



Weight Loss Agents Prior Authorization with Quantity Limit Program Summary

This program applies to FlexRx Closed, FlexRx Open, FocusRx, GenRx Closed, GenRx Open, and KeyRx.

This is a FlexRx Standard and GenRx Standard program.

POLICY REVIEW CYCLE

Effective Date
12/1/2023

Date of Origin
7/1/2013

FDA APPROVED INDICATIONS AND DOSAGE

Agent(s)	FDA Indication(s)	Notes	Ref#
Adipex-P®, Lomaira™, Phentermine* ~ Tablet Capsule	Short-term (a few weeks) adjunct in a regimen of weight reduction based on exercise, behavioral modification and caloric restriction in the management of exogenous obesity in patients with an initial BMI greater than or equal to 30 kg/m ² or greater than or equal to 27 kg/m ² in the presence of other risk factors (e.g., controlled hypertension, diabetes, hyperlipidemia).	* Generic available ~ The safety and efficacy of coadministration with other weight loss drug products, including prescribed drugs, over-the-counter preparations, and herbal products have not been established. Therefore, coadministration is not recommended.	5,6,11
Benzphetamine*~ Tablet	Short-term (a few weeks) adjunct in a regimen of weight reduction based on caloric restriction in the management of exogenous obesity for patients with an initial body mass index greater than or equal to 30 kg/m ² who have not responded to appropriate weight reducing regimen (diet and/or exercise) alone.	* Generic available ~ The safety and efficacy of coadministration with other weight loss drug products, including prescribed drugs, over-the-counter preparations, and herbal products have not been established. Therefore, coadministration is not recommended.	2
Contrave® (naltrexone/bupropion)~ Tablet ER	Adjunct to a reduced-calorie diet and increased physical activity for chronic weight management in adults with an initial body mass index (BMI) of: <ul style="list-style-type: none"> • Greater than or equal to 30 kg/m² (obese), or • Greater than or equal to 27 kg/m² (overweight) in the presence of at least one weight-related comorbidity (e.g., hypertension, type 2 diabetes mellitus, or dyslipidemia) Limitations of Use:	~ The safety and efficacy of coadministration with other weight loss drug products, including prescribed drugs, over-the-counter preparations, and herbal products have not been established. Therefore, coadministration is not recommended.	3

Agent(s)	FDA Indication(s)	Notes	Ref#
	<ul style="list-style-type: none"> Effect on cardiovascular morbidity and mortality has not been established 		
Diethylpropion* Tablet Tablet ER	<p>Short-term (a few weeks) adjunct in a regimen of weight reduction based on caloric restriction in the management of exogenous obesity for patients with an initial body mass index (BMI) greater than or equal to 30 kg/m² and who have not responded to appropriate weight reducing regimen (diet and/or exercise) alone.</p> <p>Indicated for use as monotherapy only.</p>	* Generic available	9
Phendimetrazine* Capsule ER	<p>Short-term (a few weeks) adjunct in a regimen of weight reduction based on caloric restriction in the management of exogenous obesity for patients with an initial BMI greater than or equal to 30 kg/m² or greater than or equal to 27 kg/m² in the presence of other risk factors (e.g., controlled hypertension, diabetes, hyperlipidemia) who have not responded to appropriate weight reducing regimen alone (diet and/or exercise)</p> <p>Indicated for use as monotherapy only.</p>	* Generic available	7
Phendimetrazine* Tablet	<p>Short-term (a few weeks) adjunct in a regimen of weight reduction based on caloric restriction in the management of exogenous obesity for patients with an initial BMI greater than or equal to 30 kg/m² who have not responded to appropriate weight reducing regimen alone (diet and/or exercise).</p> <p>Indicated for use as monotherapy only.</p>	* Generic available	10
Qsymia® (phentermine/topiramate)~ Capsule	<p>Adjunct to a reduced-calorie diet and increased physical activity for chronic weight management in:</p> <ul style="list-style-type: none"> Adults with an initial BMI of: <ul style="list-style-type: none"> Greater than or equal to 30 kg/m² (obese) Greater than or equal to 27 kg/m² (overweight) in the presence of at least one weight-related comorbidity such as hypertension, type 2 diabetes mellitus, or dyslipidemia Pediatric patients aged 12 years and older with BMI in the 95th percentile or greater standardized for age and sex <p>Limitations of Use:</p> <ul style="list-style-type: none"> Effect on cardiovascular morbidity and mortality has not been established 	~ The safety and efficacy of coadministration with other weight loss drug products, including prescribed drugs, over-the-counter preparations, and herbal products have not been established. Therefore, coadministration is not recommended.	1
Saxenda® (liraglutide)~ Injection solution	<p>Adjunct to a reduced-calorie diet and increased physical activity for chronic weight management in:</p> <ul style="list-style-type: none"> Adults with an initial body mass index (BMI) of: <ul style="list-style-type: none"> Greater than or equal to 30 kg/m² (obese), or Greater than or equal to 27 kg/m² (overweight) in the presence of at least one weight-related comorbidity (e.g., hypertension, type 2 diabetes mellitus, or dyslipidemia) Pediatric patients aged 12 years or older with: <ul style="list-style-type: none"> Body weight greater than 60 kg, and 	~ The safety and efficacy of coadministration with other weight loss drug products, including prescribed drugs, over-the-counter preparations, and herbal products have not been established. Therefore, coadministration is not recommended.	8

Agent(s)	FDA Indication(s)	Notes	Ref#
	<ul style="list-style-type: none"> ○ An initial BMI corresponding to greater than 30 kg/m² for adults (obese) by international cut-offs (Cole Criteria) <p>Limitations of Use:</p> <ul style="list-style-type: none"> • Safety and effectiveness in pediatric patients with type 2 diabetes have not been established • Should not be used in combination with any other GLP-1 receptor agonist 		
Wegovy® (semaglutide) ~ Subcutaneous injection solution	<p>Adjunct to a reduced calorie diet and increased physical activity for chronic weight management in:</p> <ul style="list-style-type: none"> • Adults with an initial body mass index (BMI) of: <ul style="list-style-type: none"> ○ Greater than or equal to 30 kg/m² (obese), or ○ Greater than or equal to 27 kg/m² (overweight) in the presence of at least one weight-related comorbidity (e.g., hypertension, type 2 diabetes mellitus, or dyslipidemia) • Pediatric patients aged 12 years and older with an initial BMI at the 95th percentile or greater standardized for age and sex (obesity) <p>Limitations of Use:</p> <ul style="list-style-type: none"> • Should not be coadministered with other semaglutide-containing products or with any other GLP-1 receptor agonist • Wegovy has not been studied in patients with a history of pancreatitis 	~ The safety and efficacy of coadministration with other weight loss drug products, including prescribed drugs, over-the-counter preparations, and herbal products have not been established. Therefore, coadministration is not recommended.	20
Xenical®, Orlistat Capsule*	Obesity management including weight loss and weight maintenance when used in conjunction with a reduced-calorie diet and to reduce the risk for weight regain after prior weight loss. It is indicated for obese patients with an initial body mass index (BMI) greater than or equal to 30 kg/m ² or greater than or equal to 27 kg/m ² in the presence of other risk factors (e.g., hypertension, diabetes, dyslipidemia)		4,21

See package insert for FDA prescribing information: <https://dailymed.nlm.nih.gov/dailymed/index.cfm>

CLINICAL RATIONALE

Obesity	<p>Obesity rates have increased sharply over the last 30 years, creating a global public health crisis. The National Health and Nutrition Examination Surveys show that nearly 2 of 3 US adults are overweight or obese, and 1 of 3 adults are obese. Adults with body mass index (BMI) 25-29.9 kg/m² are considered overweight; those with BMI greater than or equal to 30 kg/m² are considered obese.(14) Weight loss is difficult for most people and weight loss medications help reinforce behavioral strategies to lose weight. Medications for weight loss do not work on their own. Numerous guidelines recommend the addition of weight loss medications only in conjunction with lifestyle and behavioral modifications.(13,14,15,22)</p> <p>The American Association of Clinical Endocrinologists and American College of Endocrinology comprehensive clinical practice guidelines for medical care of patients with obesity recommends the following:(14)</p>
---------	---

- The principal outcome and therapeutic target in the treatment of obesity should be to improve the health of the patient by preventing or treating weight related complications using weight loss, not the loss of body weight per se
 - For overweight (BMI 25-29.9 kg/m²) or obese (BMI greater than or equal to 30 kg/m²) patients, evaluate for adiposity related complications:
 - Metabolic syndrome
 - Prediabetes
 - Type 2 diabetes (T2DM)
 - Dyslipidemia
 - Hypertension
 - Cardiovascular disease
 - Non-alcoholic fatty liver disease
 - Polycystic ovary syndrome
 - Female infertility
 - Male hypogonadism
 - Obstructive sleep apnea
 - Asthma/reactive airway disease
 - Osteoarthritis
 - Urinary stress incontinence
 - Gastroesophageal reflux disease
 - Depression
 - Pharmaceutical therapy should only be used as adjunct to lifestyle modifications and depends on the staging of obesity:
 - Overweight Stage 0 (BMI 25-29.9 kg/m² or 23-24.9 kg/m² in certain ethnicities* with no complications)
 - Lifestyle therapy – reduced-calorie healthy meal plan/physical activity/behavioral interventions
 - Obesity Stage 0 (BMI greater than or equal to 30 kg/m² or greater than or equal to 25 kg/m² in certain ethnicities* with no complications)
 - Lifestyle therapy – reduced-calorie healthy meal plan/physical activity/behavioral intervention
 - Weight loss medications – consider if lifestyle therapy fails to prevent progressive weight gain (BMI greater than or equal to 27 kg/m²)
 - Obesity Stage 1 (BMI greater than or equal to 25 kg/m² or greater than or equal to 23 kg/m² in certain ethnicities* with greater than or equal to 1 mild/moderate complications)
 - Lifestyle therapy – reduced-calorie healthy meal plan/physical activity/behavioral interventions
 - Weight loss medications – consider if lifestyle therapy fails to achieve therapeutic target or initiate concurrently with lifestyle therapy (BMI greater than or equal to 27 kg/m²)
 - Obesity Stage 2 (BMI greater than or equal to 25 kg/m² or greater than or equal to 23 kg/m² in certain ethnicities* with greater than or equal to 1 severe complications):
 - Lifestyle therapy – reduced-calorie healthy meal plan/physical activity/behavioral interventions
 - Weight loss medication – initiate concurrently with lifestyle therapy (BMI greater than or equal to 27 kg/m²)
 - Consider bariatric surgery (BMI greater than or equal to 35 kg/m²)
- *Certain ethnicities (A BMI cutoff point value of greater than or equal to 23 kg/m² should be used in the screening and confirmation of excess adiposity in South Asian, Southeast Asian, and East Asian adults)

The Endocrine Society clinical practice guidelines suggests medications approved for chronic weight management can be useful adjuncts to lifestyle change for patients who have been unsuccessful with diet and exercise alone. They recommend adherence to American Heart Association Guidelines (2013) [see below] which include advice for

assessment and treatment with diet and exercise, as well as bariatric surgery for appropriate candidates.(13)

- Diet, exercise, and behavioral modification should be included in all overweight and obesity management approaches for BMI greater than or equal to 25 kg/m² and other tools [e.g., pharmacotherapy (if BMI greater than or equal to 27 kg/m² with comorbidity or BMI greater than 30 kg/m²) and bariatric surgery (BMI greater than or equal to 35 kg/m² with comorbidity or BMI greater than 40 kg/m²)] should be used as adjuncts to behavioral modification to reduce food intake and increase physical activity when possible. Patients who have a history of being unable to successfully lose and maintain weight and who meet label indications are candidates for weight loss medications.
- Assessment of efficacy and safety of prescribed weight loss medications should be performed at least monthly for the first 3 months, then at least every 3 months thereafter.
- Clinicians are recommended to perform annual and symptom-based screening for major obesity related chronic conditions in all adult patients with a BMI greater than or equal to 30 kg/m², including diabetes, cardiovascular disease, hypertension, hyperlipidemia, obstructive sleep apnea, non-alcoholic fatty liver disease, osteoarthritis, and major depression.
- Prescribers should identify chronic medications, for concomitant medical conditions, that contribute to weight gain, and prescribe drugs that are weight neutral or that will promote weight loss when possible.
- If a patient's response to a weight loss medication is deemed effective (weight loss greater than or equal to 5% of body weight at 3 months) and safe, it is recommended that the medication be continued. If deemed ineffective (weight loss less than 5% at 3 months) or if there are safety or tolerability issues at any time, the medication should be discontinued and alternative medications or referral for alternative treatment approaches instead considered.
- Given the wide clinical prescribing of phentermine for greater than 20 years and lack of evidence of serious side effects, even in the absence of long-term controlled safety and efficacy data, it seems reasonable for clinicians to prescribe phentermine long term as long as the patient: 1) has no evidence of serious cardiovascular disease; 2) does not have serious psychiatric disease or a history of substance abuse; 3) has been informed about weight loss medications that are FDA approved for long-term use and told that these have been documented to be safe and effective whereas phentermine has not; 4) does not demonstrate a clinically significant increase in pulse or BP when taking phentermine; and 5) demonstrates a significant weight loss while using the medication. These aspects of care should be documented in the patient's medical record, and the off-label nature of the prescribing should be documented at each visit. Medication should be started at 7.5 or 15 mg/day initially and only increased if the patient is not achieving clinically significant weight loss. Patients should be followed at least monthly during dose escalation and then at least every 3 months when on a stable dose.

The American Heart Association/American College of Cardiology/Obesity Society Guideline (2013) suggests if weight and lifestyle history indicates the patient has never participated in a comprehensive lifestyle intervention program as defined in the guidelines (i.e., trained interventionist or nutritional professional supervision of diet, exercise, and behavior therapy), it is recommended that the patient undertake such a program before addition of adjunctive therapies (e.g., pharmacotherapy), since a substantial proportion of patients will lose sufficient weight to improve health with comprehensive lifestyle management alone. If a patient has been unable to lose weight or sustain weight loss with comprehensive lifestyle intervention and has BMI greater than or equal to 30 kg/m² or greater than or equal to 27 kg/m² with greater than or equal to 1 obesity-associated comorbid condition(s), adjunctive therapy may be considered. The expert panel did not review comprehensive evidence on pharmacotherapy for weight loss. Medications should be FDA approved and clinicians should be knowledgeable about the product label. The provider should weigh potential risks of the medication vs. potential benefits of successful weight loss for the

	<p>individual patient. If the patient is currently taking an obesity medication but has not lost at least 5% of initial body weight after 12 weeks on a maximal dose of the medication, the provider should reassess the risk-to-benefit ratio of that medication for the patient and consider discontinuation of that drug.(15)</p> <p>The Veterans Affairs and Department of Defense (VA/DoD) Clinical Practice Guideline (2020), suggests offering prescribed pharmacotherapy in patients with a BMI greater than or equal to 30 kg/m² or greater than or equal to 27 kg/m² with greater than or equal to 1 obesity-associated comorbid condition(s), in conjunction with a comprehensive lifestyle intervention.(18)</p> <p>Four centrally-acting noradrenergic agents (phentermine, diethylpropion, phendimetrazine, benzphetamine) are FDA-approved for the “short-term” (usually considered less than or equal to 12 weeks) management of obesity. However, the short-term designation was given since all were approved before the necessity of long-term treatment for obesity was established.(12) Since then some of these agents, such as phentermine and diethylpropion, have had literature published in support of long-term use.(13,19) Given the wide clinical prescribing of phentermine for greater than 20 years and lack of evidence of serious side effects, even in the absence of long-term controlled safety and efficacy data, it seems reasonable for clinicians to prescribe phentermine long term. (13) A clinical study found that diethylpropion plus a standard dietary intervention produced sustained and clinically significant weight loss over 1 year, and demonstrated safety under the cardiovascular and psychiatric point of view.(19)</p>
Pediatric Obesity	<p>Pediatric obesity has become an epidemic and international problem. In the United States, the prevalence of obesity in children has risen from 5% in 1970 to 17% in 2004. Genetics and environment are the underlying causes of the increase in pediatric obesity. Obese children and adolescents are at risk of developing the same comorbid conditions as obese and overweight adults. Obesity and overweight in children are defined on percentages specific for age and gender defined BMI values. The American Academy of Pediatrics (AAP) define obesity as a BMI greater than or equal to 95th percentile or a BMI greater than or equal to 30 kg/m², whichever is lower, and overweight as a BMI within 85th to 94th percentile for children and adolescents 2 years of age and older.(17,23)</p> <p>The AAP recommends that clinicians should assess medical and behavioral risks in any child with a BMI above the 85th percentile before initiating any intervention.(17,23) The Endocrine Society Pediatric Obesity Treatment Guidelines also recommend that clinicians should evaluate for potential comorbidities in children and adolescents with a BMI greater than or equal to 85th percentile.(16)</p> <p>The 2023 AAP guidelines recommend the use of weight loss agents in conjunction with lifestyle and behavioral changes. Pediatricians and other primary healthcare providers should treat children and adolescents for overweight with comorbidities (BMI greater than or equal to 85th percentile; comorbidities such as dyslipidemia, prediabetes, Type 2 diabetes, fatty liver disease, hypertension) and obesity (BMI greater than or equal to 95th percentile).(23)</p> <p>The 2017 Endocrine Society guidelines only recommend the use of FDA approved pharmacotherapy in pediatric patients as adjunctive therapy to lifestyle modifications of the highest intensity available and only by clinicians that are experienced in the use of anti-obesity agents.(16)</p> <ul style="list-style-type: none"> • Suggest pharmacotherapy in children or adolescents with obesity (greater than or equal to 95th percentile for age and gender) only after a formal program of intense lifestyle modifications has failed to limit weight gain or to ameliorate comorbidities. • Recommend against using obesity agents in children and adolescents less than 16 years of age who are overweight, but not obese, except in the context of clinical trials.

	<ul style="list-style-type: none"> • Anti-obesity agents should be discontinued, and patients reevaluated if the patient does not have a greater than 4% BMI reduction after 12 weeks at the medication's full dosage. • Discourages prescribing weigh loss medications off-label to pediatric patients less than 16 years of age.
Safety	<p>Phentermine has the following contraindications:(5,6,11)</p> <ul style="list-style-type: none"> • History of cardiovascular disease (e.g., coronary artery disease, stroke, arrhythmias, congestive heart failure, uncontrolled hypertension) • During or within 14 days following the administration of monoamine oxidase inhibitors • Hyperthyroidism • Glaucoma • Agitated states • History of drug abuse • Pregnancy • Nursing • Known hypersensitivity, or idiosyncrasy to the sympathomimetic amines <p>Benzphetamine has the following contraindications:(2)</p> <ul style="list-style-type: none"> • Patients with advanced arteriosclerosis, symptomatic cardiovascular disease, moderate to severe hypertension, hyperthyroidism, known hypersensitivity or idiosyncrasy to sympathomimetic amines, and glaucoma • Benzphetamine should not be given to patients who are in an agitated state or who have a history of drug abuse • Hypertensive crises have resulted when sympathomimetic amines have been used concomitantly or within 14 days following use of monoamine oxidase inhibitors • Benzphetamine tablets should not be used concomitantly with other CNS stimulants • Benzphetamine is contraindicated in women who are or may become pregnant <p>Phendimetrazine has the following contraindications:(7,10)</p> <ul style="list-style-type: none"> • Immediate release: <ul style="list-style-type: none"> ○ Known hypersensitivity or idiosyncratic reactions to sympathomimetics ○ Advanced arteriosclerosis, symptomatic cardiovascular disease, moderate and severe hypertension, hyperthyroidism, and glaucoma ○ Highly nervous or agitated ○ History of drug abuse ○ Use in combination with other CNS stimulants, including monoamine oxidase inhibitors • Extended release: <ul style="list-style-type: none"> ○ History of cardiovascular disease (e.g., coronary artery disease, stroke, arrhythmias, congestive heart failure, uncontrolled hypertension, pulmonary hypertension) ○ During or within 14 days following the administration of monoamine oxidase inhibitors ○ Hyperthyroidism ○ Glaucoma ○ Agitated states ○ History of drug abuse ○ Pregnancy ○ Nursing ○ Use in combination with other anorectic agents or CNS stimulants

- Known hypersensitivity or idiosyncratic reactions to sympathomimetics

Diethylpropion has the following contraindications:(9)

- Pulmonary hypertension, advanced arteriosclerosis, hyperthyroidism, known hypersensitivity or idiosyncrasy to the sympathomimetic amines, glaucoma, severe hypertension
- Agitated states
- Patients with a history of drug abuse
- Use in combination with other anorectic agents is contraindicated
- During or within 14 days following the administration of monoamine oxidase inhibitors, hypertensive crises may result

Phentermine/topiramate has the following contraindications:(1)

- Pregnancy
- Glaucoma
- Hyperthyroidism
- During or within 14 days following the administration of monoamine oxidase inhibitors
- Known hypersensitivity or idiosyncrasy to the sympathomimetic amines

Naltrexone/bupropion (NB) has the following:(3)

- Contraindications:
 - Uncontrolled hypertension
 - Seizure disorder or a history of seizures
 - Use of other bupropion-containing products (including, but not limited to, Wellbutrin, Wellbutrin SR, Wellbutrin XL, Aplenzin, and Zyban)
 - Bulimia or anorexia nervosa, which increase the risk for seizure
 - Chronic opioid or opiate agonist (e.g., methadone) or partial agonists (e.g., buprenorphine) use, or acute opiate withdrawal
 - Patients undergoing an abrupt discontinuation of alcohol, benzodiazepines, barbiturates, and antiepileptic drugs
 - Concomitant administration of monoamine oxidase inhibitors (MAOI). At least 14 days should elapse between discontinuation of MAOI and initiation of treatment with Contrave. There is an increased risk of hypertensive reactions when Contrave is used concomitantly with MAOIs. Starting Contrave in a patient treated with reversible MAOIs such as linezolid or intravenous methylene blue is also contraindicated
 - Known allergy to bupropion, naltrexone or any other component of Contrave. Anaphylactoid/anaphylactic reactions and Stevens-Johnson syndrome have been reported with bupropion
- Boxed warnings:
 - Contrave is not approved for use in the treatment of major depressive disorder or other psychiatric disorders. Contrave contains bupropion, the same active ingredient as some other antidepressant medications (including, but not limited to, Wellbutrin, Wellbutrin SR, Wellbutrin XL, and Aplenzin). Antidepressants increased the risk of suicidal thoughts and behavior in children, adolescents, and young adults in short-term trials. These trials did not show an increase in the risk of suicidal thoughts and behavior with antidepressant use in subjects over age 24; there was a reduction in risk with antidepressant use in subjects aged 65 and older. In patients of all ages who are started on Contrave, monitor closely for worsening, and for the emergence of suicidal thoughts and behaviors. Advise families and caregivers of the need for close observation and communication

with the prescriber. Contrave is not approved for use in pediatric patients.

Liraglutide has the following:(8)

- Contraindications:
 - Patients with a personal or family history of medullary thyroid carcinoma (MTC) or patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2)
 - Patients with a prior serious hypersensitivity reaction to liraglutide or to any of the product components
 - Pregnancy
- Boxed warnings:
 - Liraglutide causes dose-dependent and treatment-duration-dependent thyroid C-cell tumors at clinically relevant exposures in both genders of rats and mice. It is unknown whether Saxenda causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans, as the human relevance of liraglutide-induced rodent thyroid C-cell tumors has not been determined.
 - Saxenda is contraindicated in patients with a personal or family history of MTC and in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk of MTC with use of Saxenda and inform them of symptoms of thyroid tumors (e.g., a mass in the neck, dysphagia, dyspnea, persistent hoarseness). Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with Saxenda.

Orlistat has the following contraindications:(4)

- Pregnancy
- Chronic malabsorption syndrome
- Cholestasis
- Known hypersensitivity to Orlistat or to any component of this product

Semaglutide has the following:(20)

- Contraindications:
 - Personal or family history of medullary thyroid carcinoma (MTC) or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2)
 - A prior serious hypersensitivity to semaglutide or to any component of this product
- Boxed warnings:
 - Semaglutide causes dose-dependent and treatment-duration-dependent thyroid C-cell tumors at clinically relevant exposures in rodents. It is unknown whether Wegovy causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as human relevance of semaglutide-induced rodent thyroid C-cell tumors has not been determined.
 - Wegovy is contraindicated in patients with a personal or family history of MTC or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk of MTC with use of Wegovy and inform them of symptoms of thyroid tumors (e.g., a mass in the neck, dysphagia, dyspnea, persistent hoarseness). Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with Wegovy.

Co-Administration

	None of the FDA approved weight loss agents have approval for co-administration with another weight loss agent. New guidelines do not support the use of co-administration of weight loss pharmacological agents.(13,14,18) Use of non-approved drug combinations for obesity treatment should be limited to clinical trials, and patients should be informed when drugs are being used off label alone or in combination.(12)
--	--

REFERENCES

Number	Reference
1	Qsymia prescribing information. Vivus, Inc. June 2022.
2	Benzphetamine prescribing information. Nivagen Pharmaceuticals. January 2016.
3	Contrave prescribing information. Nalpropion Pharmaceuticals LLC. November 2021.
4	Xenical prescribing information. H2-Pharma, LLC. November 2022.
5	Adipex-P prescribing information. Teva Pharmaceuticals. September 2020.
6	Phentermine prescribing information. Aurolife Pharma LLC. January 2019.
7	Phendimetrazine ER prescribing information. Virtus Pharmaceuticals, LLC. May 2020.
8	Saxenda prescribing information. Novo Nordisk Inc. June 2022.
9	Diethylpropion prescribing information. Lannett Company, Inc. December 2019.
10	Phendimetrazine prescribing information. KVK-Tech, Inc. December 2019.
11	Lomaira prescribing information. KVK-Tech Inc. December 2018.
12	Yanovski SZ, Yanovski JA. Long-Term Drug Treatment for Obesity: A Systematic and Clinical Review. JAMA. 2014 Jan;311(1):74-86.
13	Apovian CM, Aronne LJ, Bessesen DH, et al. Pharmacological Management of Obesity: An Endocrine Society Clinical Practice Guideline. J Clin Endocrinol Metab. 2015 Feb;100(2):342-362.
14	American Association of Clinical Endocrinologists and American College of Endocrinology Comprehensive Clinical Practice Guidelines for Medical Care of Patients with Obesity. Endocr Pract. 2016 Jul;22(Suppl 3):1-203.
15	Jensen MD, Ryan DH, Apovian CM, et al. 2013 AHA/ACC/TOS Guideline for the Management of Overweight and Obesity in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. Circulation. 2014;129(25 Suppl 2):S102-S138.
16	Styne DM, Arslanian SA, Connor EL, et al. Pediatric Obesity – Assessment, Treatment, and Prevention: An Endocrine Society Clinical Practice Guideline. J Clin Endocrinol Metab. 2017 Jan;102(3):709-757.
17	Barlow SE, et al. Expert Committee Recommendations Regarding the Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity: Summary Report. Pediatrics. 2007 Dec;120(Suppl 4):S164-S192.
18	Department of Veterans Affairs and the Department of Defense Clinical Practice Guideline for the Management of Adult Overweight and Obesity – Version 3.0. 2020. Available at: https://www.healthquality.va.gov/guidelines/CD/obesity/ .
19	Cercato C, Roizenblatt VA, Leanca CC, et al. A Randomized Double-Blind Placebo-Controlled Study of the Long-Term Efficacy and Safety of Diethylpropion in the Treatment of Obese Subjects. Int J Obes (Lond). 2009 Aug;33(8):857-865.
20	Wegovy prescribing information. Novo Nordisk Inc. December 2022.
21	Orlistat prescribing information. H2-Pharma, LLC. June 2022.
22	American Gastroenterological Association (AGA) Clinical Practice Guideline on Pharmacological Interventions for Adults with Obesity. Gastroenterology. 2022 Nov;163(5):1198-1225.
23	American Academy of Pediatrics (AAP) Clinical Practice Guideline for the Evaluation and Treatment of Children and Adolescents with Obesity. Pediatrics. 2023 Jan;151(2):1-100.

POLICY AGENT SUMMARY PRIOR AUTHORIZATION

Target Brand Agent(s)	Target Generic Agent(s)	Strength	Targeted MSC	Available MSC	Final Age Limit	Preferred Status
	benzphetamine hcl tab	50 MG	M ; N ; O ; Y	Y		
	Diethylpropion HCl Tab 25 MG	25 MG	M ; N ; O ; Y	Y		
	Diethylpropion HCl Tab ER 24HR 75 MG	75 MG	M ; N ; O ; Y	M		
Saxenda	liraglutide (weight mngmt) soln pen-inj	18 MG/3ML	M ; N ; O ; Y	N		
Contrave	naltrexone hcl-bupropion hcl tab er	8-90 MG	M ; N ; O ; Y	N		
Xenical	Orlistat Cap 120 MG	120 MG	M ; N ; O ; Y	N		
	phendimetrazine tartrate cap er	105 MG	M ; N ; O ; Y	N		
	phendimetrazine tartrate tab	35 MG	M ; N ; O ; Y	Y		
Adipex-p	phentermine hcl cap	15 MG ; 30 ; 30 MG ; 37.5 MG	M ; N ; O ; Y	O ; Y		
Adipex-p ; Lomaira	phentermine hcl tab	37.5 MG ; 8 MG	M ; N ; O ; Y	N ; O ; Y		
Qsymia	phentermine hcl-topiramate cap er	11.25-69 MG ; 15-92 MG ; 3.75-23 MG ; 7.5-46 MG	M ; N ; O ; Y	N		
Wegovy	semaglutide (weight mngmt) soln auto-injector	0.25 MG/0.5ML ; 0.5 MG/0.5ML ; 1 MG/0.5ML ; 1.7 MG/0.75ML ; 2.4 MG/0.75ML	M ; N ; O ; Y	N		

POLICY AGENT SUMMARY QUANTITY LIMIT

Target Brand Agent Name(s)	Target Generic Agent Name(s)	Strength	QL Amount	Dose Form	Day Supply	Duration	Addtl QL Info	Allowed Exceptions	Targeted NDCs When Exclusions Exist
	Benzphetamine HCl Tab 25 MG		90	Tablets	30	DAYS			
	Benzphetamine HCl Tab 50 MG	50 MG	90	Tablets	30	DAYS			
	Diethylpropion HCl Tab 25 MG	25 MG	90	Tablet	30	DAYS			
	Diethylpropion HCl Tab ER 24HR 75 MG	75 MG	30	Tablets	30	DAYS			
	Phendimetrazine Tartrate Cap ER 24HR 105 MG	105 MG	30	Capsules	30	DAYS			
	Phendimetrazine Tartrate Tab 35 MG	35 MG	180	Tablets	30	DAYS			
	Phentermine HCl Cap 15 MG	15 MG	30	Capsules	30	DAYS			
	Phentermine HCl Cap 30 MG	30 ; 30 MG	30	Capsules	30	DAYS			
Adipex-p	Phentermine HCl Cap 37.5 MG	37.5 MG	30	Capsules	30	DAYS			

Target Brand Agent Name(s)	Target Generic Agent Name(s)	Strength	QL Amount	Dose Form	Day Supply	Duration	Addtl QL Info	Allowed Exceptions	Targeted NDCs When Exclusions Exist
Adipex-p	Phentermine HCl Tab 37.5 MG	37.5 MG	30	Tablets	30	DAYS			
Contrave	Naltrexone HCl-Bupropion HCl Tab ER 12HR 8-90 MG	8-90 MG	120	Tablets	30	DAYS			
Lomaira	Phentermine HCl Tab 8 MG	8 MG	90	Tablets	30	DAYS			
Qsymia	Phentermine HCl-Topiramate Cap ER 24HR 11.25-69 MG	11.25-69 MG	30	Capsules	30	DAYS			
Qsymia	Phentermine HCl-Topiramate Cap ER 24HR 15-92 MG	15-92 MG	30	Capsules	30	DAYS			
Qsymia	Phentermine HCl-Topiramate Cap ER 24HR 3.75-23 MG	3.75-23 MG	30	Capsules	30	DAYS			
Qsymia	Phentermine HCl-Topiramate Cap ER 24HR 7.5-46 MG	7.5-46 MG	30	Capsules	30	DAYS			
Saxenda	Liraglutide (Weight Mngmt) Soln Pen-Inj 18 MG/3ML (6 MG/ML)	18 MG/3ML	15	mLs	30	DAYS			
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	0.25 MG/0.5 ML	8	Pens	180	DAYS	* - This strength is not approvable for maintenance dosing		
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	0.5 MG/0.5 ML	8	Pens	180	DAYS	* - This strength is not approvable for maintenance dosing		
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	1 MG/0.5 ML	8	Pens	180	DAYS	* - This strength is not approvable for maintenance dosing		
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	1.7 MG/0.75 ML	4	Pens	28	DAYS			
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	2.4 MG/0.75 ML	4	Pens	28	DAYS			
Xenical	Orlistat Cap 120 MG	120 MG	90	Capsules	30	DAYS			

CLIENT SUMMARY – PRIOR AUTHORIZATION

Target Brand Agent Name(s)	Target Generic Agent Name(s)	Strength	Client Formulary
	benzphetamine hcl tab	50 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
	Diethylpropion HCl Tab 25 MG	25 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx

Target Brand Agent Name(s)	Target Generic Agent Name(s)	Strength	Client Formulary
	Diethylpropion HCl Tab ER 24HR 75 MG	75 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
	phendimetrazine tartrate cap er	105 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
	phendimetrazine tartrate tab	35 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Adipex-p	phentermine hcl cap	15 MG ; 30 ; 30 MG ; 37.5 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Adipex-p ; Lomaira	phentermine hcl tab	37.5 MG ; 8 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Contrave	naltrexone hcl-bupropion hcl tab er	8-90 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Qsymia	phentermine hcl-topiramate cap er	11.25-69 MG ; 15-92 MG ; 3.75-23 MG ; 7.5-46 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Saxenda	liraglutide (weight mngmt) soln pen-inj	18 MG/3ML	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Wegovy	semaglutide (weight mngmt) soln auto-injector	0.25 MG/0.5ML ; 0.5 MG/0.5ML ; 1 MG/0.5ML ; 1.7 MG/0.75ML ; 2.4 MG/0.75ML	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Xenical	Orlistat Cap 120 MG	120 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx

CLIENT SUMMARY – QUANTITY LIMITS

Target Brand Agent Name(s)	Target Generic Agent Name(s)	Strength	Client Formulary
	Benzphetamine HCl Tab 25 MG		FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
	Benzphetamine HCl Tab 50 MG	50 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
	Diethylpropion HCl Tab 25 MG	25 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
	Diethylpropion HCl Tab ER 24HR 75 MG	75 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
	Phendimetrazine Tartrate Cap ER 24HR 105 MG	105 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
	Phendimetrazine Tartrate Tab 35 MG	35 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
	Phentermine HCl Cap 15 MG	15 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx

Target Brand Agent Name(s)	Target Generic Agent Name(s)	Strength	Client Formulary
			Closed ; GenRx Open ; KeyRx
	Phentermine HCl Cap 30 MG	30 ; 30 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Adipex-p	Phentermine HCl Cap 37.5 MG	37.5 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Adipex-p	Phentermine HCl Tab 37.5 MG	37.5 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Contrave	Naltrexone HCl-Bupropion HCl Tab ER 12HR 8-90 MG	8-90 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Lomaira	Phentermine HCl Tab 8 MG	8 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Qsymia	Phentermine HCl-Topiramate Cap ER 24HR 11.25-69 MG	11.25-69 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Qsymia	Phentermine HCl-Topiramate Cap ER 24HR 15-92 MG	15-92 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Qsymia	Phentermine HCl-Topiramate Cap ER 24HR 3.75-23 MG	3.75-23 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Qsymia	Phentermine HCl-Topiramate Cap ER 24HR 7.5-46 MG	7.5-46 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Saxenda	Liraglutide (Weight Mngmt) Soln Pen-Inj 18 MG/3ML (6 MG/ML)	18 MG/3ML	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	1 MG/0.5ML	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	0.5 MG/0.5ML	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	1.7 MG/0.75ML	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	2.4 MG/0.75ML	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Wegovy	Semaglutide (Weight Mngmt) Soln Auto-Injector	0.25 MG/0.5ML	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx
Xenical	Orlistat Cap 120 MG	120 MG	FlexRx Closed ; FlexRx Open ; FocusRx ; GenRx Closed ; GenRx Open ; KeyRx

PRIOR AUTHORIZATION CLINICAL CRITERIA FOR APPROVAL

Module	Clinical Criteria for Approval
	<p>Initial Evaluation</p> <p>(Patient new to therapy, new to Prime, or attempting a repeat weight loss course of therapy)</p> <p>Target Agent(s) will be approved when ALL the following are met:</p> <ol style="list-style-type: none"> 1. ONE of the following: <ol style="list-style-type: none"> A. The patient is 17 years of age or over and ALL of the following: <ol style="list-style-type: none"> 1. ONE of the following: <ol style="list-style-type: none"> A. The patient has a diagnosis of obesity, confirmed by a BMI greater than or equal to 30 kg/m² OR a BMI greater than or equal to 25 kg/m² if the patient is of South Asian, Southeast Asian, or East Asian descent OR B. The patient has a BMI greater than or equal to 27 kg/m² with at least one weight-related comorbidity/risk factor/complication (e.g., diabetes, dyslipidemia, coronary artery disease) AND 2. The patient has been on a weight loss regimen of a low-calorie diet, increased physical activity, and behavioral modifications for a minimum of 6 months prior to initiating therapy with the requested agent AND 3. The patient did not achieve a weight loss of 1 pound or more per week while on the weight loss regimen prior to initiating therapy with the requested agent AND 4. The patient is currently on and will continue a weight loss regimen of a low-calorie diet, increased physical activity, and behavioral modifications OR B. The patient is 12 to 16 years of age and ALL of the following: <ol style="list-style-type: none"> 1. ONE of the following: <ol style="list-style-type: none"> A. The patient has a diagnosis of obesity, confirmed by a BMI greater than or equal to 95th percentile for age and gender OR B. The patient has a diagnosis of obesity, confirmed by a BMI greater than or equal to 30 kg/m² OR C. The patient has a BMI greater than or equal to 85th percentile for age and gender AND at least one severe weight-related comorbidity/risk factor/complication AND 2. The patient has been on a weight loss regimen of a low-calorie diet, increased physical activity, and behavioral modifications for a minimum of 6 months prior to initiating therapy with the requested agent AND 3. The patient did not achieve a weight loss of 1 pound or more per week while on the weight loss regimen prior to initiating therapy with the requested agent AND 4. The patient is currently on and will continue a weight loss regimen of a low-calorie diet, increased physical activity, and behavioral modifications AND 2. If the patient has an FDA approved indication, ONE of the following: <ol style="list-style-type: none"> A. The patient's age is within FDA labeling for the requested indication for the requested agent OR B. The prescriber has provided information in support of using the requested agent for the patient's age for the requested indication AND 3. The patient does NOT have any FDA labeled contraindications to the requested agent AND 4. The patient will NOT be using the requested agent in combination with another targeted weight loss agent for the requested indication AND 5. ONE of the following: <ol style="list-style-type: none"> A. The patient has not tried a targeted weight loss agent in the past 12 months OR B. The patient has tried a targeted weight loss agent for a previous course of therapy in the past 12 months AND the prescriber anticipates success with repeating therapy AND 6. ONE of the following: <ol style="list-style-type: none"> A. The requested agent is benzphetamine, diethylpropion, phendimetrazine, or phentermine OR

Module	Clinical Criteria for Approval
	<p>B. The requested agent is Qsymia and ONE of the following:</p> <ol style="list-style-type: none"> 1. The requested dose is 3.75mg/23mg OR 2. The patient is currently being treated with Qsymia, the requested dose is greater than 3.75 mg/23 mg AND ONE of the following: <ol style="list-style-type: none"> A. ONE of the following: <ol style="list-style-type: none"> 1. For adults, the patient has demonstrated and maintained a weight loss of greater than or equal to 5% from baseline (prior to initiation of the requested agent) OR 2. For pediatric patients aged 12 years and older, the patient has experienced a reduction of at least 5% of baseline BMI (prior to initiation of the requested agent) OR B. The patient received less than 14 weeks of therapy OR C. The patient's dose is being titrated upward OR D. The patient has received less than 12 weeks (3 months) of therapy on the 15mg/92mg strength OR 3. The prescriber has provided information in support of therapy for the requested dose for this patient OR <p>C. The requested agent is Contrave and ONE of the following</p> <ol style="list-style-type: none"> 1. The patient is newly starting therapy OR 2. The patient is currently being treated and has received less than 16 weeks (4 months) of therapy OR 3. The patient has achieved and maintained a weight loss of greater than or equal to 5% from baseline (prior to initiation of requested agent) OR <p>D. The requested agent is Xenical (orlistat) and ONE of the following:</p> <ol style="list-style-type: none"> 1. The patient is 12 to 16 years of age and ONE of the following: <ol style="list-style-type: none"> A. The patient is newly starting therapy OR B. The patient is currently being treated and has received less than 12 weeks (3 months) of therapy OR C. The patient has achieved and maintained a weight loss of greater than 4% from baseline (prior to initiation of requested agent) OR 2. The patient is 17 years of age or over and ONE of the following: <ol style="list-style-type: none"> A. The patient is newly starting therapy OR B. The patient is currently being treated and has received less than 12 weeks (3 months) of therapy OR C. The patient has achieved and maintained a weight loss of greater than or equal to 5% from baseline (prior to initiation of requested agent) OR <p>E. The requested agent is Saxenda and ALL of the following:</p> <ol style="list-style-type: none"> 1. The patient will NOT be using the requested agent in combination with another GLP-1 receptor agonist agent AND 2. ONE of the following: <ol style="list-style-type: none"> A. The patient is 18 years of age or over and ONE of the following: <ol style="list-style-type: none"> 1. The patient is newly starting therapy OR 2. The patient is currently being treated and has received less than 16 weeks (4 months) of therapy OR 3. The patient has achieved and maintained a weight loss of greater than or equal to 4% from baseline (prior to initiation of requested agent) OR B. The patient is pediatric (12 to less than 18 years of age) and BOTH of the following: <ol style="list-style-type: none"> 1. The requested agent is NOT being used to treat type 2 diabetes AND 2. ONE of the following: <ol style="list-style-type: none"> A. The patient is newly starting therapy OR B. The patient is currently being treated and has received less than 20 weeks (5 months) of therapy OR C. The patient has achieved and maintained a reduction in BMI of greater than or equal to 1% from baseline (prior to initiation of requested agent) OR <p>F. The requested agent is Wegovy and ALL of the following:</p>

Module	Clinical Criteria for Approval
	<ol style="list-style-type: none"> 1. The patient will NOT be using the requested agent in combination with another GLP-1 receptor agonist agent AND 2. The patient does NOT have a history of pancreatitis AND 3. ONE of the following: <ol style="list-style-type: none"> A. The patient is newly starting therapy OR B. The patient is currently being treated and has received less than 52 weeks (1 year) of therapy OR C. ONE of the following: <ol style="list-style-type: none"> 1. The patient is an adult AND has achieved and maintained a weight loss of greater than or equal to 5% from baseline (prior to initiation of the requested agent) OR 2. The patient is pediatric (12 to less than 18 years of age) AND has achieved and maintained a reduction in BMI of at least 5% from baseline (prior to initiation of the requested agent) <p>Length of Approval:</p> <ul style="list-style-type: none"> • For Wegovy: 12 months • For Saxenda pediatric patients (age 12 to less than 18): 5 months • For Saxenda (adults) and Contrave: 4 months • For all other agents: 3 months <p>NOTE: If Quantity Limit applies, please refer to Quantity Limit Criteria.</p> <p>Renewal Evaluation</p> <p>(Patient continuing a current weight loss course of therapy)</p> <p>Target Agent(s) will be approved when ALL of the following are met:</p> <ol style="list-style-type: none"> 1. The patient has been previously approved for the requested agent through the plan's Prior Authorization process AND 2. The patient is currently on and will continue to be on a weight loss regimen of a low-calorie diet, increased physical activity, and behavioral modifications AND 3. The patient does NOT have any FDA labeled contraindications to the requested agent AND 4. For Saxenda only, BOTH of the following: <ol style="list-style-type: none"> A. The requested agent is NOT being used to treat type 2 diabetes in pediatric patients (12 to less than 18 years of age) AND B. The patient will NOT be using the requested agent in combination with another GLP-1 receptor agonist agent AND 5. For Wegovy only, ALL of the following: <ol style="list-style-type: none"> A. The requested dose is 1.7 mg or 2.4 mg AND B. The patient will NOT be using the requested agent in combination with another GLP-1 receptor agonist agent AND C. The patient does NOT have a history of pancreatitis AND 6. The patient meets ONE of the following: <ol style="list-style-type: none"> A. The patient has achieved and maintained a weight loss greater than or equal to 5% from baseline (prior to initiation of requested agent) OR B. For Saxenda only, ONE of the following: <ol style="list-style-type: none"> 1. If the patient is 18 years of age or over, the patient has achieved and maintained a weight loss greater than or equal to 4% from baseline (prior to initiation of requested agent) OR 2. If the patient is pediatric (12 to less than 18 years of age), the patient has achieved and maintained a reduction in BMI of greater than or equal to 1% from baseline (prior to initiation of requested agent) OR C. For Qsymia only, ONE of the following:

Module	Clinical Criteria for Approval
	<ol style="list-style-type: none"> 1. For pediatric patients aged 12 years and older, the patient has achieved and maintained a reduction of at least 5% of baseline (prior to initiation of the requested agent) BMI OR 2. The patient has achieved and maintained a weight loss less than 5% from baseline (prior to initiation of requested agent) for adults, or a reduction in BMI less than 5% from baseline (prior to initiation of the requested agent) for pediatric patients aged 12 years or older, AND BOTH of the following: <ol style="list-style-type: none"> A. The patient's dose is being titrated upward (for the 3.75 mg/23 mg, 7.5 mg/46 mg or 11.25 mg/69 mg strengths only) AND B. The patient has received less than 12 weeks of therapy on the 15mg/92mg strength OR D. For Xenical (orlistat) only, ONE of the following: <ol style="list-style-type: none"> 1. The patient 12 to 16 years of age AND has achieved and maintained a weight loss greater than 4% from baseline (prior to initiation of requested agent) OR 2. The patient is 17 years of age or over AND has achieved and maintained a weight loss greater than or equal to 5% from baseline (prior to initiation of requested agent) OR E. For Wegovy only, ONE of the following: <ol style="list-style-type: none"> 1. The patient is 12 years of age and over AND has received less than 52 weeks of therapy on the maximum-tolerated dose (1.7 mg or 2.4 mg) OR 2. The patient is pediatric (12 to less than 18 years of age) AND has achieved and maintained a reduction in BMI of at least 5% from baseline (prior to initiation of the requested agent) AND 7. If the patient is 12 to less than 18 years of age, the current BMI is greater than 85th percentile for age and gender AND 8. The patient will NOT be using the requested agent in combination with another targeted weight loss agent for the requested indication <p>Length of Approval:</p> <ul style="list-style-type: none"> • Qsymia: greater than or equal to 5% weight loss from baseline (adults); greater than or equal to 5% reduction in BMI from baseline (pediatrics): 12 months • Qsymia less than 5% weight loss from baseline (adults); less than 5% reduction in BMI from baseline (pediatrics): 3 months • All other agents: 12 months <p>NOTE: If Quantity Limit applies, please refer to Quantity Limit Criteria.</p>

QUANTITY LIMIT CLINICAL CRITERIA FOR APPROVAL

Module	Clinical Criteria for Approval
	<p>Target Agent(s) will be approved when ONE of the following is met:</p> <ol style="list-style-type: none"> 1. The requested quantity (dose) does NOT exceed the program quantity limit OR 2. ALL of the following: <ol style="list-style-type: none"> A. The requested quantity (dose) exceeds the program quantity limit AND B. The requested quantity (dose) does NOT exceed the maximum FDA labeled dose for the requested indication AND C. The requested quantity (dose) cannot be achieved with a lower quantity of a higher strength that does NOT exceed the program quantity limit OR 3. ALL of the following: <ol style="list-style-type: none"> A. The requested quantity (dose) exceeds the program quantity limit AND B. The requested quantity (dose) exceeds the maximum FDA labeled dose for the requested indication AND C. The prescriber has provided information in support of therapy with a higher dose for the requested indication

Module	Clinical Criteria for Approval
	<p>Length of Approval:</p> <ul style="list-style-type: none"> • Initial Approval: <ul style="list-style-type: none"> ○ For Wegovy: 12 months ○ For Saxenda pediatric patients (age 12 to less than 18): 5 months ○ For Saxenda (adults) and Contrave: 4 months ○ For all other agents: 3 months • Renewal Approval: <ul style="list-style-type: none"> ○ Qsymia: greater than or equal to 5% weight loss from baseline (adults); greater than or equal to 5% reduction in BMI from baseline (pediatrics): 12 months ○ Qsymia. less than 5% weight loss from baseline (adults); less than 5% reduction in BMI from baseline (pediatrics): 3 months ○ All other agents: 12 months